




# Soil Monitoring Moisture Networks in Canada











- Two soil moisture monitoring networks 60X60 km resolution
- Sampling at 5, 20 and 50cm depths




- Temporal Frequency : Hourly
- Variables Observed:
   Soil temperature
   Soil Moisture
   Precipitation












### Development of Paired Soil Moisture Sampling Networks for Satellite and Model Validation

How Many Sensors?




# How representative are the network measurements compared to field conditions?






## Impact of Land-use Practices: Tillage




#### Impacts of Landuse Practices:




#### Difference in Mean Volumetric Water Content in Tilled and Non-Tilled Soils Over Time




#### Difference in the Coefficient of Variation for Volumetric Water Content in Tilled and Non-Tilled Soils Over Time





## Impact of crop cover on mean soil moisture and variability







#### Soil Moisture and Variability Time Series






Controls on soil moisture variability in Alberta. Correlation of physical processes to the first Principal Component (47% of variability) for the Alberta Drought Monitoring Network and





## Validation of Soil Moisture Estimate from Microwave Brightness Temperatures



### Correlations of satellite derived soil moisture to network observations

	LPRM C/X-Band		NSIDC X-Band		SSM/I	
ONT	R	Anomaly RMSE (%)	R	Anomaly RMSE (%)	R	Anomaly RMSE (%)
All	0.54	0.91	-0.06	1.25	0.44	1.01
MJ	0.80	0.99	0.57	1.53	0.54	1.20
JA	0.79	0.77	0.31	0.93	0.63	0.82
so	0.70	1.00	-0.57	1.25	0.25	1.05
SASK						
All	0.08	1.34	-0.04	1.38	0.02	1.54
MJ	0.71	1.65	-0.28	1.28	0.29	1.98
JA	-0.18	1.57	0.82	1.16	0.11	1.67
SO	0.23	1.27	0.11	1.81	0.02	1.62


Table 7. Relationship between in-situ and satellite derived soil moisture for 2007 and 2008 growing seasons. In-situ soil moisture is the weekly average of all available soil moisture stations within each study area. The LPRM soil moisture is derived from C-band with X-band substituted where RFI is detected, the NSIDC is derived from X-band only. Anomalies represent standard normal deviates. Values in bold are significant at p<0.05.


#### Improving soil moisture retrievals of the LPRM through parameter adjustment

	Measure	Jul 27 2007 - Oct 31 2007	May 1 2008 - Jun 23 2008	Aug 21 2008 - Oct 28 2008	Overall		
Asco.9	Bias	0.0535	-0.0174	0.0275	0.0172		
	RMSE	0.0621	0.0484	0.0394	0.0486		
Asc	d	0.9832	0.9904	0.9920	0.9893		
	R	0.2024	0.2307	0.7328	0.3261		
Desch.9	Bias	0.0741	-0.0016	0.0406	0.0330		
	RMSE	0.0833	0.0374	0.0597	0.0595		
	d	0.9723	0.9947	0.9827	0.9850		
	R	0.2500	0.6669	0.4769	0.3976		
٥,0	Bias	0.1025	0.0080	0.0839	0.0608		
W. Asa	RMSE	0.1110	0.0723	0.0890	0.0892		
LPRM-Asc	d	0.9559	0.9808	0.9680	0.9699		
	R	-0.0318	0.1741	0.7075	0.1301		
J. P. Desc	Bias	0.1181	0.0159	0.0808	0.0659		
	RMSE	0.1352	0.0854	0.1335	0.1189		
562	d	0.9389	0.9744	0.9287	0.9484		
4,	R	0.1965	0.3318	-0.0505	0.0238		
Note: $d = degree of agreement$ , $R = pairwise correlation coefficient$ .							


## Use of Network data for evaluation of data assimilation approaches

	Measure	Overall
LPRINDESG.9	Bias	0.0173
20%	RMSE	0.0483
	d	0.9894
3,	R	0.3279
7 Olas Potadi	Bias	0.0347
' Dega	RMSE	0.0531
3.5%	d	0.9880
9	R	0.1714
<del>ر</del> کو	Bias	0.0105
a drillated	RMSE	0.0373
ر المنظم	$\mathrm{d}$	0.9935
¥**	R	0.5618





## Canadian SMOS Validation Campaign 2010



- Time: May-June 2010
- Locations: Boreal Forest and Agricultural Regions of Saskatchewan
- Support: NSERC (~500K), Environment Canada in kind contributions
- Ground Support: Nested monitoring networks and BERMS; vegetation and soil moisture measurement teams
  - Aircraft Support: Environment Canada (Anne Walker)

